

Onderbouwend	Stof	Effect	Code
SPC Incivo	atorvastatine + telaprevir	↑ atorvastatine AUC 7.88 en Cmax 10.6; geCI	(1A)
Lee JE. Antimicrob Agents Chemother 2011;55:4569-74. Epub 2011 Aug 8.	atorvastatine + telaprevir	atorvastatine: ↑ Cmax van 3.5 naar 38.5 ng/ml (10.6x) en AUC van 33.8 naar 277 ng.h/ml (7.88x), t1/2 van 9.4 naar 6.7 h This was an open-label, single sequence, nonrandomized study involving 21 healthy male and female volunteers. A coformulation of 5 mg amlodipine and 20 mg atorvastatin was administered on day 1. Telaprevir was taken with food as a 750-mg dose every 8 h from day 11 until day 26, and a single dose of the amlodipine-atorvastatin combination was readministered on day 17.	

Opmerkingen

WFG 27-3-2012: beoordeling interacties HCV-middelen gedelegeerd naar de tabel die is opgesteld door prof. dr. D.M. Burger, Apotheek/Klinische Farmacie, UMC St Radboud Nijmegen. Er wordt dus ook geen alfanumerieke code (voor bewijs en potentiëel effect) meer aan toegekend.