

Filgotinib + Famotidine	MFB 7796
-------------------------	----------

Onderbouwend	Stof	Effect	Code
--			

Overig	Stof	Effect
Anderson K. Clin Pharmacol Drug Dev 2019;8:585-94. doi: 10.1002/cpdd.659.	filgotinib + famotidine	↓Cmax filgotinib met 18% (but the 90%CI remained within the 70% to 143% no-effect boundary); geen effect op AUC (GLSM ratios for AUC close to 100% and 90%CI's within the no-effect boundary); ↑Tmax 1→2 uur, geen effect op t1/2; geen effect op AUC of Cmax GS-829845; ↑Tmax 4→6 uur. Regime: filgotinib maleaat 200 mg 1-malig dag 1, 4-dagen washout; famotidine 40 mg 2dd op dag 6-9; famotidine 40 mg 2dd plus filgotinib 200 mg op dag 10 (2 uur na de ochtenddosering famotidine); studie met 13 vrijwilligers.
SPC Jyseleca	filgotinib + famotidine	geen wijziging door famotidine, geen dosisaanpassing nodig.

Opmerkingen

PubMed: verder geen hits.

Risicofactoren	
Mitigerende factoren	

	Interactie	Actie	Datum
Beslissing WG IA	Nee	Nee	21 oktober 2021